

# 3-Point block backward differentiation formula with an of-step point for the solutions of stiff chemical reaction problems

*by* Abdullah Abdullah

---

**Submission date:** 08-Nov-2022 12:57PM (UTC+0700)

**Submission ID:** 1947985182

**File name:** Paper.pdf (1.66M)

**Word count:** 7792

**Character count:** 37054



## 3-Point block backward differentiation formula with an off-step point for the solutions of stiff chemical reaction problems

Hira Soomro<sup>1</sup> · Nooraini Zainuddin<sup>1</sup> · Hanita Daud<sup>1</sup> · Joshua Sunday<sup>2</sup> · Noraini Jamaludin<sup>1</sup> · Abdullah Abdullah<sup>3</sup> · Apriyanto Mulono<sup>4</sup> · Evizal Abdul Kadir<sup>5</sup>

Received: 16 February 2022 / Accepted: 31 August 2022  
© The Author(s) 2022

### Abstract

A major challenge in simulating chemical reaction processes is integrating the stiff systems of Ordinary Differential Equations (ODEs) describing the chemical reactions due to stiffness. Thus, it would be of interest to search systematically for stiff solvers that are close to optimal for such problems. This paper presents an implicit 3-Point Block Backward Differentiation Formula with one off-step point (3POBBDF) for the solutions of first-order stiff chemical reaction problems. In deriving the method, the Lagrange polynomial was adopted as the basis function. The paper further analyses the basic properties of the 3POBBDF which include order of accuracy, consistence, zero-stability, and convergence. The stability region as well as the interval of instability of the method was also computed. To demonstrate the accuracy of the proposed approach, some famous stiff chemical reaction problems such as Robertson problem and Chemical AKZO were solved, and the results obtained were compared with those of some existing methods. The results obtained clearly show that the 3POBBDF performs better than the existing methods with which we compared our results.

**Keywords** Stiff chemical reactions · Linear multistep methods · Block off-step point · Backward differential formula · ODEs

---

✉ Hira Soomro  
soomro\_19001048@utp.edu.my; hirasoomro47@gmail.com

✉ Nooraini Zainuddin  
aini\_zainuddin@utp.edu.my

Extended author information available on the last page of the article

## 1 Introduction

Most chemical reaction problems have been known to be stiff in nature, little wonder they have attracted the attention of mathematicians recently. According to [1, 2], chemical reactions in alive species, atmospheric phenomena, mechanics and molecular dynamics, and chemical kinetics, extensively describe the stiff systems. If the time constants fluctuate significantly in magnitude, then the system is known as stiff. The time constants here refer to the rates of decay of perturbations [3]. If a system diverges considerably in terms of time constants, we must use smaller time steps to deal with expeditiously decaying terms.

In this paper, an implicit 3-Point Block Backward Differentiation Formula with one off-step point (3POBBDF) is derived for the solutions of stiff chemical reaction problems of the form,

$$y'(x) = f(x, y), y(a) = y_0, x \in [a, b] \quad (1.1)$$

where  $f$  is assumed to be continuous and satisfies the existence and uniqueness theorem.

For Ordinary Differential Equations (ODEs) such as non-linear or linear, stiff or non-stiff ODEs, different numerical methods have been developed [4–6]. It must be noted that employing the wrong method for a model can result in a slow or incorrect solution [7]. Problems with the form (1.1) are generally divided into two categories. The first category is a non-stiff ODEs, in which explicit methods are employed together with certain error control. Another category is stiff ODEs. [8] was the first to use the term "stiff" in a scenario requiring chemical kinetics. The only way to find answers to stiff problems is to employ implicit methods because explicit methods are slow and sometimes fail to deliver an accurate solution [9]. The Backward Differentiation Formula (BDF) is one of the well-known classes of implicit methods for finding the solution of stiff ODEs. Developing a mathematical model for stiff ODEs considers a few factors such as selection of step size, stability, accuracy, and computational cost [10, 11].

Stiffness is defined in different ways in literature as it does not have a specific definition. Equation (1.1) is said to be stiff according to Lambert [12] if the Jacobian matrix  $\frac{\partial f}{\partial y}$  for the eigenvalues  $\lambda_t(x)$  satisfies these conditions:

$$Re(\lambda_t(x)) < 0, t = 1, 2, \dots, n$$

$$\max_t |Re(\lambda_t(x))| \gg \min_t |Re(\lambda_t(x))|$$

where  $\lambda_t$  are the eigenvalues and the ratio  $\frac{\max_{t=1,2,\dots,n} |Re(\lambda_t)|}{\min_{t=1,2,\dots,n} |Re(\lambda_t)|}$  is called stiffness ratio.

Many studies have been directed into the improvement of multistep block methods for the solution of (1.1). Several studies have recently focused on improving the performance of different forms of multistep block methods encompassing estimated solution accuracy and computational time; see [10, 13–19]. While several block methods (based on Adams formulas) were proposed previously emphasizing

ODEs of non-stiff systems, the challenge has always been how to find an appropriate implicit method for solving stiff ODEs that have A-stability properties. Since block methods' most serious flaw appears to be their ability to sustain stability [20], it is necessary to develop algorithms with suitable stability properties.

Gear [21] initially introduced the BDF, which is known to be efficient in handling complex stiff ODE problems. This BDF approach has been extended in papers such as [15, 22–25]. Single-step and multi-step block techniques are the two types of block methods. Rosser [26] introduced the standard single-step method known as the Runge–Kutta method. [27] explored the single-step implicit A-stable r-block approach. [28] developed a fourth-order block approach for multistep methods that can be used as a predictor–corrector method. [29] proposed a Modified extended BDF method for numerical solutions of stiff ODEs. [15] devised a method for solving first-order ODEs using 2-point and 3-point block BDF. [30] developed the 5th order 2-Point Block Backward Differentiation Formula to improve the existing BBDF. Authors in articles [31–33] also improved the BBDF method by extending the method and adding future points. Many researchers have explored other block methods such as [34] who developed a seven-step block method for the solution of first order IVPs in ODEs and [35] considered the approximate solutions to third-order boundary value problems (BVPs) of ODEs.

Despite having many advantages, the block method also possessed a major setback which pointed out that the order of interpolation points must not exceed the order of the differential equations [36, 37]. Because of this setback, addition of off-step points in the block were introduced and named as hybrid method. Hybrid methods are highly efficient and have been reported to circumvent the “Dahlquist Zero-Stability Barrier” condition by introducing function evaluation at off-step points which takes some time in its development but provides better approximation than two conventional methods (Runge–Kutta and linear multistep methods) [13, 36]. [38] proposed function evaluation at an off-step and named as “Hybrid” method. [9] derived higher-order method with more than one off-step point. [39] were the first who introduce BDF methods with some off-step points. The hybrid Obrechko BDF methods with some off-step points were presented in [40]. Hybrid Block Second Derivative Backward Differentiation Formula (HBSDBDF) is proposed by [41] which provides simultaneous solutions of IVPs in each block without any predictor formula. [42] formulated Block Backward Differentiation Formula with two off-step points (BBDFO(6)). [23] have studied block methods with various off-step points. [43–45] have described a closely related methods that involves off-step points. In modifying the selection of the most acceptable off-step points, a similar method is used from [46, 47].

Motivated from the above literature, this study is aimed at developing a 3-Point Block Backward Differentiation Formula with one off-step point (3POBBDF) using Lagrange polynomials as basis function to solve chemical reactions problems comprising stiff first order ODEs. The method will be employed in solving stiff chemical reaction problems of the form (1.1). Several points have been examined for the selection of off-step points and hence it was concluded that selecting the points where the step is halved will lead to zero-stable formulae. Compared to the methods developed by Khalsaraei and Molayi [48], and Ismail [49], the advantage of the

proposed method is that the solutions are approximated with off-step points concurrently and provide better accuracy. The organization of the paper is as follow; Section 2 briefly describes the formulation of the method. Analysis of convergence properties is examined in Section 3. Section 4 elaborates the implementation of the derived method. List of tested problems with results are presented in section 5. Section 6 includes discussions of the results. Graphical representation of the results is displayed in section 7. consists of Numerical problems with their results respectively. Furthermore, Sections 8 describes the conclusions.

## 2 Derivation of the 3POBBDF

This section consists of a detailed explanation of how the proposed 3POBBDF for solving (1.1) is formulated. The two starting values,  $x_{n-1}$  and  $x_n$  of identical step size which is fixed as  $h = x_{n+1} - x_n$  and one off-step point at  $x_{n+\frac{5}{2}}$  is considered, as shown in Fig. 1.

The interval  $[a, b]$  in (1.1) is split into a sequence of blocks in a 3-point block method (see Fig. 1). For finding out the solution of (1.1),  $y_{n-1}$  and  $y_n$  are used as the values of previous block to evaluate the future points at  $y_{n+1}, y_{n+2}, y_{n+\frac{5}{2}}$  and  $y_{n+3}$  with constant step size simultaneously. The off-step point  $y_{n+\frac{5}{2}}$  is added to all three formulas to increase the accuracy. The 3POBBDF we shall derive is a  $k$ -step Linear Multi-step Method (LMM) defined according to [12] as,

$$\sum_{j=0}^k \alpha_j y_{n+j} = h \sum_{j=0}^k \beta_j f_{n+j} \quad (2.1)$$

$h$  represents the step size,  $\alpha_j$  and  $\beta_j$  are the unknown constants that needs to be decided and  $k$  indicates the number of steps of the method. Presuming that  $\alpha_j \neq 0$ , and both  $\alpha_0$  and  $\beta_0$  are not equal to zero. The LMM (2.1) is extended by adding an off-step point to generate a new form of LMM given by,




Fig. 1 3-Point Block Backward Differentiation Formula with one off-step point

$$\sum_{j=0}^4 \alpha_{j-1} y_{n+(j-1)} + \alpha_{\frac{5}{2}} y_{\frac{5}{2}} = h\beta_{T,T} f_{n+T}, \text{ for } T = 1, 2, \frac{5}{2}, 3 \quad (2.2)$$

The function  $f_{n+T}$  in Eq. (2.2) is evaluated by using Lagrange interpolation polynomial ( $P(x)$ ) with interpolation points such as  $(x_{n-1}, y_{n-1}), (x_n, y_n), (x_{n+1}, y_{n+1}), (x_{n+2}, y_{n+2}), (x_{n+\frac{5}{2}}, y_{n+\frac{5}{2}})$  and  $(x_{n+3}, y_{n+3})$ . The process is given as,

$$P(x) = \sum_{j=0}^4 L_j(x) f(x_{n+3-j}), \quad (2.3)$$

whereas,

$$L_j = \prod_{\substack{i=0 \\ i \neq j}}^4 \frac{x - x_{n+3-i}}{x_{n+3-j} - x_{n+3-i}} \text{ for } i = 0, \frac{1}{2}, 1, 2, 3, 4$$

Expanding the polynomial  $L_j$  by substituting  $s = \frac{x-x_{n+3}}{h}$  implying  $x = sh + x_{n+3}$ . The interpolating polynomial is further differentiated with respect to  $s$  provides the coefficient values of  $\alpha$  and  $\beta$  as presented in Table 1.

Furthermore, (2.2) is shaped up after substituting all values taken from Table 1. Hence, the corresponding formulas for the points  $y_{n+1}, y_{n+2}, y_{n+\frac{5}{2}}$ , and  $y_{n+3}$  take the following form:

$$\begin{aligned} y_{n+1} &= \frac{3}{56}y_{n-1} - \frac{3}{5}y_n + 3y_{n+2} - \frac{64}{35}y_{n+\frac{5}{2}} + \frac{3}{8}y_{n+3} - \frac{3}{2}hf_{n+1}, \\ y_{n+2} &= -\frac{1}{98}y_{n-1} + \frac{3}{35}y_n - \frac{3}{7}y_{n+1} + \frac{384}{245}y_{n+\frac{5}{2}} - \frac{3}{14}y_{n+3} - \frac{6}{7}hf_{n+2}, \\ y_{n+\frac{5}{2}} &= -\frac{75}{9088}y_{n-1} + \frac{147}{2272}y_n - \frac{1225}{4544}y_{n+1} + \frac{3675}{2272}y_{n+2} - \frac{3675}{9088} \\ y_{n+3} + \frac{105}{142}hf_{n+\frac{5}{2}} \quad y_{n+3} &= \frac{3}{343}y_{n-1} - \frac{16}{245}y_n + \frac{12}{49}y_{n+1} - \frac{48}{49} \\ y_{n+2} + \frac{3072}{1715}y_{n+\frac{5}{2}} + \frac{12}{49}hf_{n+3}. \end{aligned} \quad (2.4)$$

Equation (2.4) is the 3POBBDF for the solution of stiff chemical reaction problems of the form (1.1).

**Table 1** Coefficients of 3POBBDF

	$\alpha_{-1}$	$\alpha_0$	$\alpha_1$	$\alpha_2$	$\alpha_{\frac{5}{2}}$	$\alpha_3$	$\beta_{T,T}$
$y_{n+1}$	$-\frac{3}{56}$	$\frac{3}{5}$	1	-3	$\frac{64}{35}$	$-\frac{3}{8}$	$\frac{3}{2}$
$y_{n+2}$	$\frac{1}{98}$	$-\frac{3}{35}$	$\frac{3}{7}$	1	$-\frac{384}{245}$	$\frac{3}{14}$	$\frac{6}{7}$
$y_{n+\frac{5}{2}}$	$\frac{75}{9088}$	$-\frac{147}{2272}$	$\frac{1225}{4544}$	$-\frac{3675}{2272}$	1	$\frac{3675}{9088}$	$-\frac{105}{142}$
$y_{n+3}$	$-\frac{3}{343}$	$\frac{16}{245}$	$-\frac{12}{49}$	$\frac{48}{49}$	$-\frac{3072}{1715}$	1	$-\frac{12}{49}$

### 3 Analysis of basic properties of the 3POBBDF

The following section comprises the analysis of basic properties of the proposed method. These properties include order, consistency, zero-stability and convergence. The region of absolute stability of the method shall also be determined.

#### 3.1 Order and error constant of the method

**Definition 3.1** (*Order and Error Constant*)

The LMM (2.1) associated with the linear difference operator is said to be of order  $p$  if  $C_1 = C_2 = \dots = C_p = 0$  and  $C_{p+1} \neq 0$ . The term  $C_{p+1} \neq 0$  is called the error constant of the method

According to [12] and [50], the local truncation error related with (2.1), for determining the order and error constant of the proposed method is well-defined by the linear difference operator  $L(y(x))$  as,

$$L(y(x)) = \sum_{j=0}^{k+s-1} (a_{j-1}y(x+Th) - h\beta_{T,T}y(x+Th)) \quad (3.1)$$

The function  $y(x+Th)$  with its derivative  $y'(x+Th)$  are expanded and collecting terms in (3.1) will be obtained such as,

$$L[y(x);h] = C_0y(x) + C_1y'(x) + \dots + C_qh^qy^{(q)}(x) + \dots \quad (3.2)$$

On the application of Eq. (3.2) on 3POBBDF derived in Eq. (2.4) using the coefficients presented in Table 1, we obtain

$$\begin{aligned} C_0 &= \sum_{j=0}^{k+s-1} \alpha_{j-(k-1),T} = [0000]^T, C_1 = \sum_{j=0}^{k+s-1} t\alpha_{j-(k-1),T} - \beta_{T,T} = [0000]^T, \\ C_2 &= \sum_{j=0}^{k+s-1} \frac{t^2}{2!} \alpha_{j-(k-1),T} - T\beta_{T,T} \\ &= [0000]^T, \therefore C_5 = \sum_{j=0}^{k+s-1} \frac{t^5}{5!} \alpha_{j-(k-1),T} - \frac{(T)^4}{4!} \beta_{T,T} = [0000]^T, \\ C_6 &= \sum_{j=0}^{k+s-1} \frac{t^6}{6!} \alpha_{j-(k-1),T} - \frac{(T)^5}{5!} \beta_{T,T} = \left[ \frac{1}{80} - \frac{1}{280} \frac{21}{6250} - \frac{1}{245} \right]^T \neq [0000]^T \end{aligned} \quad (3.3)$$

As a result, the error constant of the proposed method is obtained as

$$C_6 = \begin{bmatrix} \frac{1}{80} \\ -\frac{280}{21} \\ \frac{6250}{1} \\ -\frac{1}{245} \end{bmatrix}.$$

According to Definition 3.1,  $c_0 = c_1 = \dots = c_5 = [00000]^T$ . The term  $c_6$  represents the error constant. Therefore, this can be demonstrated that the proposed method is of order five.

### 3.1.1 Consistency of the method

#### Definition 3.2 (Consistency)

The LMM (2.1) is said to be consistent if it is of order  $p \geq 1$ .

It is therefore clear that the 3POBBDF is consistent since it has order  $p=5$  which satisfies Definition 3.2

### 3.1.2 Zero-stability of the method

#### Definition 3.3 (Zero-stable)

If no root of the characteristic polynomial has a modulus greater than one and every root with modulus one is simple, then LMM (2.2) is said to be zero-stable

To be a convergent method, zero-stability is one of the main criteria which will be discussed here. [13] proposed the scalar test to determine the stability of the method (2.4) as

$$y' = \lambda y, \quad \lambda < 0, \quad (3.4)$$

where  $\lambda$  represents the complex constant with  $\text{Re}(\lambda) < 0$ . Equation (3.4) is substituted in Eq. (2.4), therefore it precedes as,

$$y_{n+1} = \frac{3}{56}y_{n-1} - \frac{3}{5}y_n + 3y_{n+2} - \frac{64}{35}y_{n+\frac{5}{2}} + \frac{3}{8}y_{n+3} - \frac{3}{2}h\lambda y_{n+1}, \quad (3.5)$$

$$y_{n+2} = -\frac{1}{98}y_{n-1} + \frac{3}{35}y_n - \frac{3}{7}y_{n+1} + \frac{384}{245}y_{n+\frac{5}{2}} - \frac{3}{14}y_{n+3} - \frac{6}{7}h\lambda y_{n+2},$$

$$y_{n+\frac{5}{2}} = -\frac{75}{9088}y_{n-1} + \frac{147}{2272}y_n - \frac{1225}{4544}y_{n+1} + \frac{3675}{2272}y_{n+2} - \frac{3675}{9088}y_{n+3} + \frac{105}{142}h\lambda y_{n+\frac{5}{2}},$$

$$y_{n+3} = \frac{3}{343}y_{n-1} - \frac{16}{245}y_n + \frac{12}{49}y_{n+1} - \frac{48}{49}y_{n+2} + \frac{3072}{1715}y_{n+\frac{5}{2}} + \frac{12}{49}h\lambda y_{n+3}$$

The matrix is then created by inscribing Eq. (3.5) into the matrix form as follows:



$$\begin{bmatrix} -\frac{3}{8} & \frac{64}{35} & -3 & 1 + \frac{3}{2}h\lambda \\ \frac{3}{3675} & -\frac{245}{384} & 1 + \frac{6}{7}h\lambda & \frac{3}{7} \\ \frac{14}{9088} & 1 - \frac{142}{3072}h\lambda & -\frac{3675}{48} & \frac{1225}{4544} \\ 1 - \frac{12}{49}h\lambda & -\frac{1715}{1715} & \frac{2272}{49} & -\frac{12}{49} \end{bmatrix} \begin{bmatrix} y_{n+3} \\ y_{n+\frac{5}{2}} \\ y_{n+2} \\ y_{n+1} \end{bmatrix} = \begin{bmatrix} -\frac{3}{5} & 0 & \frac{3}{56} & 0 \\ \frac{3}{35} & 0 & -\frac{1}{98} & 0 \\ \frac{147}{2272} & 0 & -\frac{98}{9088} & 0 \\ -\frac{16}{245} & 0 & \frac{3}{343} & 0 \end{bmatrix} \begin{bmatrix} y_n \\ y_{n-\frac{1}{2}} \\ y_{n-1} \\ y_{n-2} \end{bmatrix},$$

which is equivalent to,

$$CY_n = DY_{n-1}, \quad (3.6)$$

$C$  and  $D$  are appropriately selected  $m \times m$  matrix coefficients,  $Y_n = [y_{n+3}, y_{n+\frac{5}{2}}, y_{n+2}, y_{n+1}]^T$  and  $Y_{n-1} = [y_n, y_{n-\frac{1}{2}}, y_{n-1}, y_{n-2}]^T$ . The stability polynomial  $R(t, H)$  correlated with the method in (2.2) is defined by

$$R(t, H) = |Ct - D|,$$

$$\begin{aligned} R(t, H) = & \frac{-13977t^2}{79147250} + \frac{44523Ht^2}{158294500} + \frac{9H^2t^2}{97412} - \frac{2883852t^3}{344125} - \frac{811134Ht^3}{807625} - \frac{15070599H^2t^3}{39573625} \\ & - \frac{2889H^3t^3}{48706} + \frac{41616t^4}{48706} - \frac{179199Ht^4}{97412} + \frac{159417H^2t^4}{97412} - \frac{40851H^3t^4}{48706} + \frac{810H^4t^4}{3479}. \end{aligned} \quad (3.7)$$

Based on the stability polynomial in (3.7), the zero-stability is obtained by replacing  $H = h\lambda = 0$  into (3.7), this yield,

$$R(t, 0) = \frac{-13977t^2}{79147250} - \frac{2883852t^3}{344125} + \frac{41616t^4}{48706}. \quad (3.8)$$

Solving  $R(t, 0) = 0$ , gives the roots for  $t$ , as  $t = 0, 0, -0.000186375, 0.999767$ . Since all the roots lie within  $|t| \leq 1$  described in Definition 4.1; hence, it is concluded that the method 3POBBDF is zero-stable.

### 3.1.3 Convergence of the method

#### Definition 3.4 (Convergence)

Consistency and zero-stability are required conditions for the LMM (2.1) to be convergent

Therefore, we conclude that the newly derived 3POBBDF is convergent since it is consistent and zero-stable.

### 3.1.4 Stability region of the method

#### Definition 3.5 (A-stable)

The term "A-stable" refers to a method that is stable for all  $\lambda$  in the left-half plane ( $Re(\lambda) \leq 0$ ). The stability region for A-stable method covers up the complete negative left-half plane

In this section, the stability region of the 3POBBDF(5) presented in Eq. (2.4) is plotted by using the stability polynomial given in (3.7). The boundary of the stability region is described by the set of points defined by  $t = e^{i\theta}, 0 \leq \theta \leq 2\pi$ . The root condition ( $|t| \leq 1$ ) of the stability polynomial must be tested at multiple grid points in order to determine the boundary of the stability region. Figure 2 shows the region of absolute stability for the system in (2.4).

It is observed from this figure that the stability region lies outside of the bounded region. Since most of the region is in the left half-plane therefore, according to Definition 3.5, the method is A-stable. Thus, it can be implied that the constructed 3POBBDF method is appropriate for solving stiff problems.

### 3.1.5 Comparison of stability regions

The stability region of the proposed 3POBBDF(5) is depicted in the Fig. 3 below, as opposed to the existing stability region of the Block Backward Differentiation Formula of Order 6 (BBDFO(6)) developed by [42]. On the complex  $h\lambda$  plane, Fig. 3 illustrates the region including both methods, namely the 3POBBDF(5) and the BBDFO(6).

The interval of instability of the current 3POBBDF(5) and the BBDFO(6) [42] are then compared. The derived 3POBBDF(5) has a larger stability area, as shown in Table 2. This implies that the proposed 3POBBDF(5) is more computationally reliable than BBDFO(6) developed by [42].

## 4 Implementation and Algorithm of the 3POBBDF

This section discusses the implementation and algorithms of the proposed method.




Fig. 2 Stability region of 3POBBDF(5)




Fig. 3 Regions of Stability for 3POBBDF(5) and BBDFO(6)

#### 4.1 Implementation of the 3POBBDF

By using Newton iteration, implementation of the 3POBBDF(5) method having one off-step is briefly mentioned in the generalized form. An increment notation is introduced for stipulating the iteration. In which  $y_{n+1}^{(i+1)}$  will represent the  $(i+1)$ <sup>th</sup> iterative value of  $y_{n+1}$  [51].

The main limitation for the implementation of the 3POBBDF(5) predominantly is that it is not self-starting. The back values are therefore calculated using the Euler method in this case.

By applying Newton's iteration, (2.4) is solved and the number of iterations is limited to two. Newton's iteration for  $y_{(n+1)}^{(i+1)}$ ,  $y_{(n+2)}^{(i+1)}$ ,  $y_{(n+\frac{5}{2})}^{(i+1)}$ ,  $y_{(n+3)}^{(i+1)}$  points in the block takes the form

$$y_{(n+1)}^{(i+1)} = y_{n+1}^{(i)} - \frac{F_1 y_{n+1}^{(i)}}{F_1' y_{n+1}^{(i)}},$$

$$y_{(n+2)}^{(i+1)} = y_{n+2}^{(i)} - \frac{F_2 y_{n+2}^{(i)}}{F_2' y_{n+2}^{(i)}},$$

**Table 2** Comparison of instability region of 3POBBDF(5) with BBDFO(6)

Methods	Interval of Instability
3POBBDF(5)	(0, 3.3375)
BBDFO(6)	(0, 10.05)

$$y_{(n+\frac{5}{2})}^{(i+1)} = y_{n+\frac{5}{2}}^{(i)} - \frac{F_{\frac{5}{2}} y_{n+\frac{5}{2}}^{(i)}}{F'_{n+\frac{5}{2}} y_{n+\frac{5}{2}}^{(i)}},$$

$$y_{(n+3)}^{(i+1)} = y_{n+3}^{(i)} - \frac{F_3 y_{n+3}^{(i)}}{F'_{n+3} y_{n+3}^{(i)}}. \quad (4.1)$$

where  $i = 0, 1$ .

We need to set the formula for  $y_{n+1}$

$$y_{n+1} = \frac{3}{56} y_{n-1} - \frac{3}{5} y_n + 3y_{n+2} - \frac{64}{35} y_{n+\frac{5}{2}} + \frac{3}{8} y_{n+3} - \frac{3}{2} h f_{n+1}.$$

Let

$$F_1 = y_{n+1} - 3y_{n+2} + \frac{64}{35} y_{n+\frac{5}{2}} - \frac{3}{8} y_{n+3} + \frac{3}{2} h f_{n+1} - \psi_1,$$

where  $\psi_1$  represent the back values

$$\psi_1 = \frac{3}{56} y_{n-1} - \frac{3}{5} y_n,$$

Referring to (4.1),  $F'_1$  for  $y_{n+1}$  is also required which is,

$$F'_1 = 1 + \frac{3h}{2} \frac{\partial f_{n+1}}{\partial y_{n+1}},$$

So, the (4.1) for  $y_{n+1}^{(i+1)}$  becomes,

$$y_{n+1}^{(i+1)} - y_{n+1}^{(i)} = - \frac{(y_{n+1}^{(i)} - 3y_{n+2}^{(i)} + \frac{64}{35} y_{n+\frac{5}{2}}^{(i)} - \frac{3}{8} y_{n+3}^{(i)} + \frac{3}{2} h f_{n+1}^{(i)} - \psi_1)}{1 + \frac{3h}{2} \frac{\partial f_{n+1}}{\partial y_{n+1}}},$$

$$\left(1 + \frac{3h}{2} \frac{\partial f_{n+1}}{\partial y_{n+1}}\right) (y_{n+1}^{(i+1)} - y_{n+1}^{(i)}) = -y_{n+1}^{(i)} + 3y_{n+2}^{(i)} - \frac{64}{35} y_{n+\frac{5}{2}}^{(i)} + \frac{3}{8} y_{n+3}^{(i)} - \frac{3}{2} h f_{n+1}^{(i)} + \psi_1, \quad (4.2)$$

Similarly, Newton iteration for  $y_{n+2}, y_{n+\frac{5}{2}}$  and  $y_{n+3}$

$$\left(1 + \frac{6h}{7} \frac{\partial f_{n+2}}{\partial y_{n+2}}\right) (y_{n+2}^{(i+1)} - y_{n+2}^{(i)}) = -\frac{3}{7} y_{n+1}^{(i)} - y_{n+2}^{(i)} + \frac{384}{245} y_{n+\frac{5}{2}}^{(i)} - \frac{3}{14} y_{n+3}^{(i)} - \frac{6}{7} h f_{n+2}^{(i)} + \psi_2, \quad (4.3)$$

$$\left(1 - \frac{105h}{142} \frac{\partial f_{n+\frac{5}{2}}}{\partial y_{n+\frac{5}{2}}}\right) \left(y_{n+\frac{5}{2}}^{(i+1)} - y_{n+\frac{5}{2}}^{(i)}\right) = -\frac{1225}{4544} y_{n+1}^{(i)} + \frac{3675}{2272} y_{n+2}^{(i)} - y_{n+\frac{5}{2}}^{(i)} - \frac{3675}{9088} y_{n+3}^{(i)} + \frac{105}{142} h f_{n+\frac{5}{2}}^{(i)} + \psi_{\frac{5}{2}} \quad (4.4)$$

$$\left(1 - \frac{12h}{49} \frac{\partial f_{n+3}}{\partial y_{n+3}}\right) \left(y_{n+3}^{(i+1)} - y_{n+3}^{(i)}\right) = \frac{12}{49} y_{n+1}^{(i)} - \frac{48}{49} y_{n+2}^{(i)} + \frac{3072}{1715} y_{n+\frac{5}{2}}^{(i)} - y_{n+3}^{(i)} + \frac{12}{49} h f_{n+3}^{(i)} + \psi_3 \quad (4.5)$$

Let  $e_{n+1}^{(i+1)} = y_{n+1}^{(i+1)} - y_{n+1}^{(i)}$  be the increment value from  $i^{\text{th}}$  to  $(i+1)^{\text{th}}$  iterations. Therefore, Eqs. (4.2)-(4.5) are then represented in matrix form as

$$\begin{bmatrix} 1 + \frac{3h}{2} \frac{\partial f_{n+1}}{\partial y_{n+1}} & -3 & \frac{64}{35} & -\frac{3}{8} \\ \frac{3}{7} & 1 + \frac{6h}{7} \frac{\partial f_{n+2}}{\partial y_{n+2}} & -\frac{384}{245} & \frac{3}{14} \\ \frac{1225}{4544} & -\frac{3675}{2272} & 1 - \frac{105h}{142} \frac{\partial f_{n+\frac{5}{2}}}{\partial y_{n+\frac{5}{2}}} & \frac{3675}{9088} \\ -\frac{12}{49} h \lambda & \frac{48}{49} & -\frac{3072}{1715} & 1 - \frac{12}{49} \frac{\partial f_{n+3}}{\partial y_{n+3}} \end{bmatrix} \begin{bmatrix} e_{n+1}^{(i+1)} \\ e_{n+2}^{(i+1)} \\ e_{n+\frac{5}{2}}^{(i+1)} \\ e_{n+3}^{(i+1)} \end{bmatrix} = \begin{bmatrix} -1 & 3 & -\frac{64}{35} & \frac{3}{8} \\ -\frac{3}{7} & -1 & \frac{384}{245} & -\frac{3}{14} \\ -\frac{1225}{4544} & \frac{3675}{2272} & -1 & -\frac{3675}{9088} \\ \frac{12}{49} & -\frac{48}{49} & \frac{3072}{1715} & -1 \end{bmatrix} \begin{bmatrix} y_{n+1}^{(i)} \\ y_{n+2}^{(i)} \\ y_{n+\frac{5}{2}}^{(i)} \\ y_{n+3}^{(i)} \end{bmatrix} + \begin{bmatrix} -\frac{3}{2} & 0 & 0 & 0 \\ 0 & -\frac{6}{7} & 0 & 0 \\ 0 & 0 & \frac{105}{142} & 0 \\ 0 & 0 & 0 & \frac{12}{49} \end{bmatrix} \begin{bmatrix} f_{n+1}^{(i)} \\ f_{n+2}^{(i)} \\ f_{n+\frac{5}{2}}^{(i)} \\ f_{n+3}^{(i)} \end{bmatrix} + \begin{bmatrix} \psi_1 \\ \psi_2 \\ \psi_{\frac{5}{2}} \\ \psi_3 \end{bmatrix}$$

Consequently, we will get the approximation values of  $y_{n+1}, y_{n+2}, y_{n+\frac{5}{2}}$  and  $y_{n+3}$  as

$$y_{n+1}^{(i+1)} = y_{n+1}^{(i)} + e_{n+1}^{(i+1)},$$

$$y_{n+2}^{(i+1)} = y_{n+2}^{(i)} + e_{n+2}^{(i+1)},$$

$$y_{n+\frac{5}{2}}^{(i+1)} = y_{n+\frac{5}{2}}^{(i)} + e_{n+\frac{5}{2}}^{(i+1)},$$

and

$$y_{n+3}^{(i+1)} = y_{n+3}^{(i)} + e_{n+3}^{(i+1)}.$$

#### 4.1.1 Algorithm of the 3POBBDF in Code

This section will elaborate on how the developed code works with the proposed method. Initially, the code will begin with three points having one off-step block method. It is employed in  $PE(CE)^r$ , the type where  $P$  and  $C$  stand for predictor and corrector correspondingly. Whereas  $E$  indicates the evaluation of function  $f$ . The power  $r = 2$  shows the number of iterations as presented by [51] that is required to converge three points having one off-step block method corrector formulae. The algorithm of the proposed method in the code is given below,

Step 1:  $P$  = Predictor values  $y_{n+m}$  for  $m = 1, 2, \frac{5}{2}, 3$ ,

Step 2:  $E$  = Evaluate  $f_{n+m}$  for  $m = 1, 2, \frac{5}{2}, 3$ ,

Step 3:  $C$  = Corrector value  $y_{n+m}$  for  $m = 1, 2, \frac{5}{2}, 3$ ,

Step 4:  $E$  = Evaluate  $f_{n+m}$  for  $m = 1, 2, \frac{5}{2}, 3$ .

### 5 Test problems

To test the reliability of the proposed 3POBBDF(5) method, some numerical results are obtained by applying the method on some well-known chemical reaction problems. Comparison is done with other multistep methods. The 3POBBDF(5) method has been coded in C++.

#### Problem 5.1 (Stiff Chemical Reaction Problem)

A chemistry problem with a stiff system is considered below,

$$y_1' = -0.013y_2 - 1000y_1y_2 - 2500y_1y_3,$$

$$y_2' = -0.013y_2 - 1000y_1y_2,$$

$$y_3' = -2500y_1y_3,$$

with initial value  $y_1 = 0, y_2 = 1$  and  $y_3 = 1, 0 \leq x \leq 2$ .

This problem is solved using  $h = 10^{-5}$  for 3POBBDF(5). Table 3 shows the integration results for this problem at  $x = 2$ . Compared with the formulas of

**Table 3** Numerical results of Problem 5.1

$x$	$y_i$	Exact solution	Absolute Error in 3POBBDF(5)	Error in [48]	Error in [49]
2.0	$y_1$	0.00003616933169289	1.1E-17	0.61E-16	0.82E-10
	$y_2$	0.9815029948230	2.29E-11	0.53E-10	0.61E-05
	$y_3$	1.0184933882440	4.39E-11	0.74E-10	0.57E-05

two-step hybrid method by Khalsaraei and Molayi [48], and Ismail's method [49], the 3POBBDF(5) method provides more accurate results in terms of Absolute error.

**Problem 5.2** (*Chemical AKZO Nobel Problem*)

The chemical AKZO noble problem is a chemical process given by a stiff system with six non-linear differential equations taken from [52] is considered. Mathematically the problem is described as,

$$\frac{dy}{dx} = F(y), \quad y(0) = y_0, \quad y \in \mathbb{R}^6, \quad 0 \leq x \leq 180.$$

The function  $F(y)$  is defined by

$$\frac{dy}{dx} = \begin{pmatrix} -2r_1 + r_2 - r_3 - r_4 \\ \frac{1}{2}r_1 - r_4 - \frac{1}{2}r_5 + F_{in} \\ r_1 - r_2 + r_3 \\ -r_2 + r_3 - 2r_4 \\ -r_2 - r_3 + r_5 \\ -r_5 \end{pmatrix},$$




where  $r_i$  and  $F_{in}$  are auxiliary variables given by



$$\begin{aligned} r_1 &= k_1 y_1^4 y_2^{\frac{1}{2}}, & k_1 &= 18.7, & r_2 &= k_2 y_3 y_4 k_2 = 0.58, \\ r_3 &= \frac{k_2}{K} y_3 y_5 K = 34.4, & r_4 &= k_3 y_1 y_4^2, & k_3 &= 0.09, \\ r_5 &= k_4 y_6^2 y_2^{\frac{1}{2}}, & k_4 &= 0.42, \end{aligned}$$

$$F_{in} = k_1 A \left( \frac{p(O_2)}{H} - y_2 \right) k_1 A = 3.3 p(O_2) = 0.9 H = 737.$$

The initial vector  $y_0 = (0.437, 0.00123, 0, 0, 0, 0.367)^T$ .

*Background of the Chemical AKZO Nobel Problem:* This problem comes from AKZO Nobel Central Research in Arnhem, the Netherlands. It defines a chemical process that mixes MBT and CHA with a continuous supply of oxygen. The important consequent species is CBS. AKZO Nobel problem contains following reaction equations,





An equilibrium is explained by the last equation

$$K_S^1 = \frac{[MBT.CHA]}{[MBT].[CHA]},$$

while the velocities of others describe reactions are given by

$$r_1 = k_1[MBT]^4.[O_2]^{\frac{1}{2}},$$

$$r_2 = k_2[MBTS].[CHA],$$

$$r_3 = \frac{k_2}{K}[MBT].[CBS],$$

$$r_4 = k_3[MBT].[CHA]^2,$$

$$r_5 = k_4[MBT.CHA]^2.[O_2]^{\frac{1}{2}},$$

respectively. Here the concentration is denoted by square brackets '[.]'.  $F_{in}$  indicated the inflow of oxygen per unit volume and satisfies a

$$F_{in} = k_l A \left( \frac{p(O_2)}{H} - [O_2] \right),$$

where  $k_l A$  is the coefficient of mass transfer,  $H$  represents Henry constant, and the partial oxygen pressure is denoted by  $p(O_2)$ .  $p(O_2)$  is supposed to be independent of  $[O_2]$ . The parameters  $k_1, k_2, k_3, k_4, K, A, p(O_2)$  and  $H$  are given constants. The procedure begins with a mixture of 0.437 mol/L [MBTS] and 0.367 mol/L [MBTS.CHA]. At the start, the oxygen concentration is 0.00123 mol/L. There is no additional presence of species at first. The simulation is completed in the time interval [0 180 min]. The mathematical formulation of the above description is simply obtained by identifying the concentrations [MBT],  $[O_2]$ , [MBTS], [CHA], [CBS], [MBT.CHA] with  $y_1, \dots, y_6$ , respectively. The numerical results by proposed method at the end point.

( $x_{end} = 180$ ) are shown in Table 4.

### Problem 5.3 (Robertson Problem)

A stiff system of three non-linear ODEs is involved in the Robertson problem. In 1966, Robertson proposed it [53]. [9] came up with the name ROBER. The file



rober.f contained the software part of the problem which can be found in [54]. The problem is described mathematically in the form.

$$\frac{dy}{dx} = F(y), \quad y(0) = y_0,$$

with  $y \in \mathbb{R}^3$ ,  $x \in [0, X]$  and the function  $F$  is specified by  $(f_1, f_2, f_3)^T$  where,




$$f_1 = -0.04y_1 + 10^4 y_2 y_3,$$

$$f_2 = 0.04y_1 - 10^4 y_2 y_3 - 3 \times 10^7 y_2^2,$$

$$f_3 = 3 \times 10^7 y_2^2,$$

with the initial value  $y_1 = 1, y_2 = 0$  and  $y_3 = 0$ .

*Origin of the Problem:* Robertson [53] defined the ROBER problem as the kinetics of an autocatalytic reaction. The composition of the reactions is as follows:



$A$ ,  $B$ , and  $C$  are the chemical species and  $k_1$ ,  $k_2$  and  $k_3$  are the rate constants. The following mathematical model, consisting of a set of three ODEs, can be put up under some ideal conditions [55] and the assumption that the mass action law is implemented to the rate functions.

**Table 4** Numerical results of Problem 5.2

$x_{end}$	$y_i$	Absolute Error in 3POBBDF(5)	Error in [48]
180	$y_1$	$6.92288 \times 10^{-6}$	$1.16162 \times 10^{-1}$
	$y_2$	$1.16287 \times 10^{-8}$	$1.11941 \times 10^{-3}$
	$y_3$	$3.55564 \times 10^{-6}$	$1.62125 \times 10^{-1}$
	$y_4$	$1.97555 \times 10^{-7}$	$3.39591 \times 10^{-3}$
	$y_5$	$1.71447 \times 10^{-5}$	$1.64618 \times 10^{-1}$
	$y_6$	$2.12229 \times 10^{-6}$	$1.98954 \times 10^{-1}$

$$\begin{pmatrix} y_1' \\ y_2' \\ y_3' \end{pmatrix} = \begin{pmatrix} -k_1 y_1 + k_3 y_2 y_3 \\ k_1 y_1 - k_2 y_2^2 - k_3 y_2 y_3 \\ k_2 y_2^2 \end{pmatrix},$$

with  $(y_1(0), y_2(0), y_3(0))^T = (y_{01}, y_{02}, y_{03})^T$ , where  $y_1, y_2$  and  $y_3$  refers to the concentrations of  $A, B$ , and  $C$  respectively, and  $y_{01}, y_{02}$  and  $y_{03}$  are concentrations at time  $t = 0$ . In the test problem,  $k_1 = 0.04, k_2 = 3 \times 10^7$  and  $k_3 = 10^4$  are the numerical values of rate constants with the initial concentrations  $y_{01} = 1, y_{02} = 0$  and  $y_{03} = 0$ .

The cause for stiffness is the substantial variation in reaction rate constants. This system features a modest, very rapid initial transient phase for problems arising from chemical kinetics. This is followed by a highly smooth fluctuation of the components, which would be acceptable for a numerical technique with a large stepsize. For  $x \in [0, 4000]$ , the ODE system is integrated. The numerical results by proposed method at the points.

$x = 0.4, 40, 4000$  are shown in Table 5.

#### Problem 5.4 (Stiff Chemical Reaction Problem)

Consider the stiff system IVPs.

$$y_1' = -1002y_1 + 1000y_2^2$$

$$y_2' = y_1 - y_2(1 + y_2),$$

with the initial conditions as  $y_1(0) = 1$  and  $y_2(0) = 1$  having exact solutions as,

$$y_1 = \exp(-2x)$$

$$y_2 = \exp(-x)$$

This problem is solved at  $x = 50$  by the new method and compared the results with forth order Two step hybrid method with one off-step point [48]. Here the stepsize  $h = 0.05$  is used for compared and proposed methods. The smaller stepsize can also be used to get more accurate results. Table 6 shows the numerical results.

**Table 5** Numerical results of Problem 5.3

$x$	$y_i$	Absolute Error in 3POBBDF(5)	Error in [48]
0.4	$y_1$	7.183E-08	9.851E-01
	$y_2$	1.227E-11	3.386E-05
	$y_3$	7.188E-08	1.479E-02
40	$y_1$	1.040E-04	7.158E-01
	$y_2$	4.010E-09	9.185E-06
	$y_3$	1.044E-04	2.841E-01
4000	$y_1$	8.395E-05	1.832E-01
	$y_2$	5.251E-10	9.942E-07
	$y_3$	8.398E-05	8.167E-01

## 6 Discussion

The results generated by the developed 3POBBDF(5) method are displayed in Tables 3, 4, 5, 6. In Table 3, comparison on the basis of absolute error was made using the 3POBBDF(5) of order 5 at the point  $x = 2$  with the methods of Khalsaraei and Molayi [48] and Ismail's method [49] at the stepsize  $h = 10^{-5}$ . In Table 4, 3POBBDF(5) method is integrated at  $h = 10^{-5}$  and the direct comparison has been made with the two step hybrid method with one off-step point [48] by taking the solution points at  $x = 180$ . Table 3 shows that the 3POBBDF(5) is more accurate and performs relatively better than the method [48] and significantly better than the method [49]. However, Table 4 shows the superiority of 3POBBDF(5) over the method [48]. Table 5 shows the numerical integration results for the problem 7.3 at  $x = 0.4, 40$  and  $4000$  using the proposed 3POBBDF(5) approach in comparison to that of [48] at  $h = 0.001$ . The comparison of the absolute errors in Table 5 for different points of  $x$  demonstrates the dominance of 3POBBDF(5) over the method given in [48] in terms of accuracy at  $h = 0.05$ . According to Table 6, it is observed that the 3POBBDF(5) method of order 5 is more effective than the two-step hybrid method of fourth order with one off-step point [48]. In Table 6, it can be seen that the 3POBBDF(5) method of order 5 performs better than the fourth order Two step hybrid method with one off-step point [48]. As shown by tables (3–6), 3POBBDF(5) is well suited to nonlinear chemical reaction problems of a stiff nature.

## 7 Graphical representations of numerical results

The tested chemical reaction problems (5.1)–(5.4) are also compared with ode15s. Figures 4, 5, 6, and 7 are the graphs demonstrating the accuracy of the 3POBBDF(5) compared with the stiff solver ode15s. It can be examined from the graphs that the results of the 3POBBDF(5) method and the results of ode15s are nearly identical at specific intervals. These figures demonstrate that the proposed method is well suited for stiff chemical reaction problems, since the solutions produced by the 3POBBDF(5) coincide with the well-known ode15s code of MATLAB.

## 8 Conclusion

In the present paper, an implicit 3POBBDF(5) method has been derived for the numerical solution of stiff systems of first-order IVPs arising from chemical reactions such as Chemical AKZO, Robertson problem, and stiff chemical problems. The

**Table 6** Numerical results of Example 5.4

$x$	$y_i$	Absolute Error in 3POBBDF(5)	Error in [48]
50	$y_1$	7.38E-24	7.14E-21
	$y_2$	4.83E-25	3.34E-19




Fig. 4 Graphs of intervals vs solution values for Problem 5.1

approach is based on the block BDF, in which at each step of the integration, three approximate solutions are generated instantaneously. Based on stability analysis of the 3POBBDF(5), the method is consistent and zero-stable; thus, the 3POBBDF(5) is convergent. Since it has an A-stability property, the method is declared suitable for solving stiff ODEs. Comparison between the stability regions was also made which shows that the 3POBBDF(5) has a smaller interval of instability than the BBDF(6) suggested by [42]. The numerical results obtained through the 3POBBDF(5) and compared to [48] and [49] evidenced that by applying the 3POBBDF(5) method,




Fig. 5 Graphs of intervals vs solution values for Problem 5.2




Fig. 6 Graphs of intervals vs solution values for Problem 5.3

the accuracy of the numerical solutions in terms of absolute error at specific points is improved. Hence, the proposed method can be successfully applied on stiff systems generated from chemical reactions because of their high order accuracy and wider stability region. Therefore, it can be concluded that the 3POBBDF(5) could be an appropriate substitute solver for stiff ODEs. To improve efficiency, further research can be implemented by using variable step sizes for solving ODEs would be beneficial.




Fig. 7 Graphs of intervals Vs solution values for Problem 5.4

**Acknowledgements** This research was funded by the National Collaborative Research Fund between Universiti Teknologi PETRONAS and Universiti Malaysia Pahang with cost centre 015MC0-036 and International Collaborative Research Fund between Universiti Teknologi PETRONAS and Universitas Islam Indragiri, Riau, Indonesia with cost centre 015ME0-220.

## Declarations

**Conflict of interest** The authors declare no conflict of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References


1. R.C. Aiken, Proceedings of the International Conference on Stiff Computation, April 12–14, 1982, Park City, Utah. Volume II, Utah univ salt lake city dept of chemical engineering (1982)
2. M. Mehdizadeh Khalsaraei, A. Shokri, M. Molayi, The new high approximation of stiff systems of first order IVPs arising from chemical reactions by k-step L-stable hybrid methods. *Iran. J. Math. Chem.* **10**(2), 181–193 (2019)
3. J. Vigo-Aguiar, H. Ramos, A new eighth-order A-stable method for solving differential systems arising in chemical reactions. *J. Math. Chem.* **40**(1), 71–83 (2006)
4. M. Sheikholeslami, Z. Shah, A. Shafee, I. Khan, I. Tlili, Uniform magnetic force impact on water based nanofluid thermal behavior in a porous enclosure with ellipse shaped obstacle. *Sci. Rep.* **9**(1), 1–11 (2019)
5. A. Hussanan, Z. Ismail, I. Khan, A.G. Hussein, S. Shafie, Unsteady boundary layer MHD free convection flow in a porous medium with constant mass diffusion and Newtonian heating. *Eur. Phys. J. Plus.* **129**(3), 1–16 (2014)
6. Y. Qin, Q. Lou, Differential transform method for the solutions to some initial value problems in chemistry. *J. Math. Chem.* **59**(4), 1046–1053 (2021)
7. M. Falati, G. Hojjati, Integration of chemical stiff ODEs using exponential propagation method. *J. Math. Chem.* **49**(10), 2210–2230 (2011)
8. C.F. Curtiss, J.O. Hirschfelder, Integration of stiff equations. *Proc. Natl. Acad. Sci. USA* **38**(3), 235 (1952)
9. G.H. Wanner, Solving ordinary differential equations II. (Springer, Berlin, 1996).
10. M.T. Chu, H. Hamilton, Parallel solution of ODE's by multiblock methods. *SIAM J. Sci. Stat. Comput.* **8**(3), 342–353 (1987)
11. L. Aceto, D. Trigiante, On the A-stable methods in the GBDF class. *Nonlinear Anal. Real World Appl.* **3**(1), 9–23 (2002)
12. J. Lambert, *Computational methods in ordinary differential equations*. New York (1973)
13. G.G. Dahlquist, A special stability problem for linear multistep methods. *BIT Numer. Math.* **3**, 27–43 (1963)
14. Z.A. Majid, M. Suleiman, F. Ismail, M. Othman, 2-Point implicit block one-step method half Gauss-Seidel for solving first order ordinary differential equations. *MATEMATIKA* **19**, 91–100 (2003)
15. Z.B. Ibrahim, K.I. Othman, M. Suleiman, Implicit r-point block backward differentiation formula for solving first-order stiff ODEs. *Appl. Math. Comput.* **186**(1), 558–565 (2007)
16. Z. A. Majid, M. Bin Suleiman, Z. Omar, 3-Point implicit block method for solving ordinary differential equations. *Bull. Malay. Math. Sci. Soc. Second Ser.* **29**(1), 23–31 (2006)

17. F. Ismail, Y.L. Ken, M. Othman, Explicit and implicit 3-point block methods for solving special second order ordinary differential equations directly. *Int. J. Math. Anal.* **3**(5), 239–254 (2009)
18. Z.B. Ibrahim, K.I. Othman, M. Suleiman, 2-Point block predictor-corrector of backward differentiation formulas for solving second order ordinary differential equations directly. *Chiang Mai J. Sci.* **39**(3), 502–510 (2012)
19. I. S. Mohd Zawawi, Z.B. Ibrahim, K.I. Othman, Derivation of diagonally implicit block backward differentiation formulas for solving stiff initial value problems. *Math. Probl. Eng.* **2015**, 1, (2015)
20. L.F. Shampine, H. Watts, Block implicit one-step methods. *Math. Comput.* **23**(108), 731–740 (1969)
21. C.W. Gear, Numerical initial value problems in ordinary differential equation. *COMM. ACM* **14**, 185–190 (1971)
22. S. Yatim, Z. Ibrahim, K. Othman, M. Suleiman, A numerical algorithm for solving stiff ordinary differential equations. *Math. Probl. Eng.* **2013**, 1 (2013)
23. N. Abasi, M. Suleiman, N. Abbasi, H. Musa, 2-Point block BDF method with off-step points for solving stiff ODEs. *J. Soft Comput. Appl.* **1**, 1–15 (2014)
24. N. Zainuddin, Z.B. Ibrahim, Block method for third order ordinary differential equations. In: *AIP Conference Proceedings*, 2017, vol. 1870, no. 1, p. 050009 (AIP Publishing LLC)
25. Z.B. Ibrahim, N. Mohd Noor, K.I. Othman, Fixed coefficient A ( $\alpha$ ) stable block backward differentiation formulas for stiff ordinary differential equations. *Symmetry* **11**(7), 46 (2019)
26. J.B. Rosser, A Runge-Kutta for all seasons. *SIAM Rev.* **9**(3), 417–452 (1967)
27. H.A. Watts, L.F. Shampine, A-stable block implicit one-step methods. *BIT Numer. Math.* **12**(2), 252–266 (1972)
28. D. Voss, S. Abbas, Block predictor-corrector schemes for the parallel solution of ODEs. *Comput. Math. Appl.* **33**(6), 65–72 (1997)
29. J. Cash, Modified extended backward differentiation formulae for the numerical solution of stiff initial value problems in ODEs and DAEs. *J. Comput. Appl. Math.* **125**(1–2), 117–130 (2000)
30. N. Nasir, Z.B. Ibrahim, K.I. Othman, M. Suleiman, Numerical solution of first order stiff ordinary differential equations using fifth order block backward differentiation formulas. *Sains Malays* **41**(4), 489–492 (2012)
31. H. Musa, M.B. Suleiman, F. Ismail, N. Senu, Z.B. Ibrahim, An improved 2-point block backward differentiation formula for solving stiff initial value problems. In: *AIP conference proceedings*, 2013, vol. 1522, no. 1, pp. 211–220: American Institute of Physics.
32. H. Musa, M. Suleiman, F. Ismail, A-stable 2-point block extended backward differentiation formula for solving stiff ordinary differential equations. In: *AIP Conference Proceedings*, 2012, vol. 1450, no. 1, pp. 254–258: American Institute of Physics.
33. H. Musa, M. Suleiman, N. Senu, Fully implicit 3-point block extended backward differentiation formula for stiff initial value problems. *Appl. Math. Sci.* **6**(85), 4211–4228 (2012)
34. S. Gebregiorgis, H. Muleta, A seven-step block multistep method for the solution of first order stiff differential equations. *Momona Ethiop. J. Sci.* **12**(1), 72–82 (2020)
35. H. Ramos, M.A. Rufai, A two-step hybrid block method with fourth derivatives for solving third-order boundary value problems. *J. Comput. Appl. Math.* **1**, 113419 (2021)
36. B.S. Kashkari, M.I. Syam, Optimization of one step block method with three hybrid points for solving first-order ordinary differential equations. *Results Phys.* **12**, 592–596 (2019)
37. G. Kumlang, S. Adee, Y. Skwame, Implicit two step Adam-Moulton hybrid block method with two offstep points for solving stiff ordinary differential equations. *J. Nat. Sc. Res.* **3**(9), 77–82 (2013)
38. W.B. Gragg, H.J. Stetter, Generalized multistep predictor-corrector methods. *J. ACM* **11**(2), 188–209 (1964)
39. M. Ebadi, M. Gokhale, Hybrid BDF methods for the numerical solutions of ordinary differential equations. *Numer. Algorithms* **55**(1), 1–17 (2010)
40. A. Shokri, A. Shokri, The hybrid Obrechhoff BDF methods for the numerical solution of first order initial value problems. *Acta Univ. Apulensis* **38**, 23–33 (2014)
41. O. Akinfenwa, R. Abdulganiy, B. Akinnukawe, S. Okunuga, Seventh order hybrid block method for solution of first order stiff systems of initial value problems. *J. Egypt. Math. Soc.* **28**(1), 1–11 (2020)
42. A.A. Nasarudin, Z.B. Ibrahim, H. Rosali, On the integration of stiff ODEs using block backward differentiation formulas of order six. *Symmetry* **12**(6), 952 (2020)
43. L.K. Yap, F. Ismail, Block methods with off-steps points for solving first order ordinary differential equations. In: *International Conference on Mathematical Sciences and Statistics 2013*, 2014, pp. 275–284. Springer.

44. H. Musa, Diagonally implicit super class of block backward differentiation formula with off-step points for solving stiff initial value problems. *Education* **6**, 1 (2016)
45. Z.O. Ra'ft Abdelrahim, J.O. Kuboye, New hybrid block method with three off-step points for solving first order ordinary differential equations. *Am. J. Appl. Sci* **13**, 209–212 (2016)
46. G. Ajileye, S. Amoo, O. Ogwumu, Hybrid block method algorithms for solution of first order initial value problems in ordinary differential equations. *J. Appl. Comput. Math.* **7**(2), 1–4 (2018)
47. P. Agarwal, I.H. Ibrahim, A new type OF hybrid multistep multiderivative formula for solving stiff IVPs. *Adv. Diff. Equ.* **2019**(1), 1–14 (2019)
48. M.M. Khalsarai, A. Shokri, M. Molayi, The new class of multistep multiderivative hybrid methods for the numerical solution of chemical stiff systems of first order IVPs. *J. Math. Chem.* **58**(9), 1987–2012 (2020)
49. G.A. Ismail, I.H. Ibrahim, New efficient second derivative multistep methods for stiff systems. *Appl. Math. Model.* **23**(4), 279–288 (1999)
50. E. Suli and D. F. Mayers, *An Introduction to Numerical Analysis* (Cambridge University Press, Cambridge, 2003)
51. N. M. N. A. K. I. O. Zarina Bibi Ibrahim, Fixed Coefficient  $A(\alpha)$  Stable Block Backward Differentiation Formulas for Stiff Ordinary Differential Equations (2019)
52. K. Eriksson, C. Johnson, A. Logg, Explicit time-stepping for stiff ODEs. *SIAM J. Sci. Comput.* **25**(4), 1142–1157 (2004)
53. H. Robertson, *The Solution of a Set of Reaction Rate Equations Numerical Analysis* (Thompson Book Co., Washington, 1967)
54. F. Mazzia, J.R. Cash, K. Soetaert, A test set for stiff initial value problem solvers in the open source software R: Package deTestSet. *J. Comput. Appl. Math.* **236**(16), 4119–4131 (2012)
55. R.C. Aiken, *Stiff computation* (Oxford University Press, Oxford, 1985)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Authors and Affiliations

Hira Soomro<sup>1</sup>  · Nooraini Zainuddin<sup>1</sup> · Hanita Daud<sup>1</sup> · Joshua Sunday<sup>2</sup> · Noraini Jamaludin<sup>1</sup> · Abdullah Abdullah<sup>3</sup> · Apriyanto Mulono<sup>4</sup> · Evizal Abdul Kadir<sup>5</sup>

Joshua Sunday  
joshuasunday2000@yahoo.com

<sup>1</sup> Mathematical and Statistical Sciences, Institute of Autonomous System, Department of Fundamental and Applied Sciences, Universiti Teknologi PETRONAS, 32610 Seri Iskandar, Perak, Malaysia

<sup>2</sup> Department of Mathematics, University of Jos, Jos 930003, Nigeria

<sup>3</sup> Department of Information System, Universitas Islam Indragiri, Tembilahan, Indonesia

<sup>4</sup> Department of Food Science, Universitas Islam Indragiri, Tembilahan, Indonesia

<sup>5</sup> Department of Informatics Engineering, Faculty of Engineering, Universitas Islam Riau, Kota Pekanbaru, Indonesia



# 3-Point block backward differentiation formula with an of-step point for the solutions of stif chemical reaction problems

---

## ORIGINALITY REPORT

---

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

14%

PUBLICATIONS

4%

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

4%

★ Mohammad Mehdizadeh Khalsaraei, Ali Shokri, Maryam Molayi. "The new class of multistep multiderivative hybrid methods for the numerical solution of chemical stiff systems of first order IVPs", *Journal of Mathematical Chemistry*, 2020

Publication

---

Exclude quotes Off

Exclude matches < 1%

Exclude bibliography On