

The Odd Beta Prime-G Family of Probability Distributions.pdf

by

Submission date: 03-Jun-2023 09:05PM (UTC+0700)

Submission ID: 2108059556

File name: The Odd Beta Prime-G Family of Probability Distributions.pdf (460.76K)

Word count: 2552

Character count: 12261

The Odd Beta Prime-G Family of Probability Distributions: Properties and Applications to Engineering and Environmental Data

Ahmad Abubakar Suleiman ^{1,2*}, Mahmud Othman ^{3,*}, ¹Aliyu Ismail Ishaq⁴, Hanita Daud ¹, Rachmah Indawati⁵, Mohd Lazim Abdullah⁶, Abdullah Husin³

¹ Fundamental and Applied Sciences Department, Universiti Teknologi PETRONAS 32610 Seri Iskandar, Perak Darul Ridzuan, Malaysia; ahmad_22000579@utp.edu.my (A.A.S.), hanita_daud@utp.edu.my (H.D.)

² Department of Statistics, Aliko Dangote University of Science and Technology, Wudil 713281, Nigeria

³ Universitas Islam Indragiri, Tembilahan, Indonesia; mahmod.othman@unisi.ac.id (M.O.), abdialam@yahoo.com (A.H.)

⁴ Department of Statistics, Ahmadu Bello University, Zaria 810107, Nigeria; aiishaq@abu.edu.ng, binishaq05@gmail.com (A.I.I.)

⁵ Faculty of Public Health, Airlangga University, Surabaya, Indonesia; rachmah.indawati@fkm.unair.ac.id (R.I.)

⁶ Management Science Research Group, Faculty of Ocean Engineering Technology, and Informatics, Universiti Malaysia Terengganu, Kuala Terengganu, Malaysia; lazim_m@umt.edu.my (M.L.A.)

* Correspondence: ahmad_22000579@utp.edu.my (A.A.S.); mahmod.othman@utp.edu.my (M.O.)

† This paper is an extended version of our paper published in Presented at the 1st International Online Conference on Mathematics and Applications; Available online: <https://iocma2023.sciforum.net/>.

Abstract: In this work, a novel generalized family of distributions called the odd beta prime is introduced. The linear representations of the proposed family are obtained. The expressions for the moments, moment generating function, and entropy are derived. A three-parameter special sub-model of the proposed family called the odd beta prime-exponential distribution is proposed. Finally, two real data sets are used to illustrate the usefulness and flexibility of the proposed distribution.

Keywords: odd beta prime generalized family; exponential distribution; T-X transformer; moments; moment generating function; entropy



Citation: Suleiman, A.A.; Othman, M.; Ishaq, A.I.; Daud, H.; Indawati, R.; Abdullah, M.L.; Husin, A. *Comput. Sci. Math. Forum* 2023, 6, x. <https://doi.org/10.3390/xxxxx>

Published: date: 28 April 2023



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

A popular research field is the construction of novel approaches for extending the existing distributions. The two interesting approaches to expanding a probability distribution are the T-X technique proposed by [1] and modified by [2]. The cumulative distribution function (cdf) for the generalizing family of distributions using this approach is given as

$$F(x) = \int_a^{W(G(x))} v(m) dm = V \{W(G(x))\}, \quad (1)$$

where $v(m)$ is the pdf of the random variable $M \in [a, b]$, such that $-\infty \leq a < b \leq \infty$

and $w(G(x))$ is a link function of any cdf of continuous distributions that take different forms. If we consider the odd function form, $w(G(x)) = \frac{G(x)}{1-G(x)}$, then the cdf of the T-X class

will be

$$F(x) = \int_0^{\frac{G(x)}{1-G(x)}} v(m) dm = V \left\{ \frac{G(x)}{1-G(x)} \right\}. \tag{2}$$

Many authors constructed extended generalized families by using the T-X approach. For example, see beta-G [3], Kw-G type-1 [4], gamma-X [5], exponentiated T-X [1], Weibull-G [6], generalized odd Lindley-G [7], and Maxwell-Weibull [8].

In this study, we consider the odd function form, $W(G(x)) = \frac{G(x)}{1-G(x)}$. Also, we

considered the beta prime distribution for $v(m) = \frac{1}{B(c,d)} \frac{x^{c-1}}{(1+x)^{c+d}}$, $x > 0$.

Therefore, we now define the odd beta prime-G family with cdf given as

$$F(x) = \frac{B_{\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}}(c,d)}{B(c,d)}, \quad x \geq 0. \tag{3}$$

The probability distribution function (pdf) of odd beta prime-G family is

$$f(x) = \frac{g(x;\varepsilon)}{B(c,d)[1-G(x;\varepsilon)]^2} \frac{\left(\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}\right)^{c-1}}{\left[1 + \left(\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}\right)\right]^{c+d}}; \quad x \in \mathfrak{R}, \tag{4}$$

where $G(x;\varepsilon)$ is a cdf of a baseline distribution with parameter ε , $g(x;\varepsilon)$ is the pdf of the baseline distribution, $c > 0$ and $d > 0$ are the shape parameters.

Here, we are motivated to propose a new flexible family of distribution called the odd beta prime generalized (OBP-G) family, which provides greater accuracy and flexibility in fitting real-life data.

This article unfolds as follows. In Section 2, linear representations of the proposed family are derived. Some statistical properties are studied and obtained in Section 3. A special sub-model of the proposed family is introduced in Section 4. In Section 5, the performance of the proposed distribution is illustrated via two applications to real data sets. Finally, Section 6 concludes the article.

2. Linear Representations

This section presents important linear representations of the OBP-G family density function defined in (4).

Let us consider the generalized Binomial expansion as follows.

$$(1+m)^{-\eta} = \sum_{i=0}^{\infty} \binom{-\eta}{i} M^i = \sum_{i=0}^{\infty} (-1)^i \binom{-\eta+i-1}{i} M^i. \tag{5}$$

Applying (5) into (4), we get

$$\begin{aligned} f(x) &= \frac{g(x;\varepsilon)}{B(c,d)[1-G(x;\varepsilon)]^2} \left(\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}\right)^{c-1} \sum_{i=0}^{\infty} (-1)^i \binom{c+d+i-1}{i} \left(\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}\right)^i, \\ &= \frac{g(x;\varepsilon)}{B(c,d)} \sum_{i=0}^{\infty} (-1)^i \binom{c+d+i-1}{i} \frac{(G(x;\varepsilon))^{c+i-1}}{(1-G(x;\varepsilon))^{c+i+1}}. \end{aligned} \tag{6}$$

Using the generalized Binomial expansion for $|z| < 1$, yields

$$(1-m)^{-n} = \sum_{j=0}^{\infty} \frac{\Gamma(n+j)}{j! \Gamma n} z^j. \tag{7}$$

Substituting (7) into (6), we obtain

$$\begin{aligned} f(x) &= \frac{g(x; \varepsilon)}{B(c, d)} \sum_{i=0}^{\infty} (-1)^i \binom{c+d+i-1}{i} (G(x; \varepsilon))^{c+i-1} \sum_{j=0}^{\infty} \frac{\Gamma(c+i+j+1)}{j! \Gamma(c+i+1)} (G(x; \varepsilon))^j, \\ &= \frac{g(x; \varepsilon)}{B(c, d)} \sum_{i,j=0}^{\infty} \Psi_{i,j} (G(x; \varepsilon))^{c+i+j-1}, \end{aligned} \tag{8}$$

where $\Psi_{i,j} = (-1)^i \binom{c+d+i-1}{i} \frac{\Gamma(c+i+j+1)}{j! \Gamma(c+i+1)}$.

3. Statistical Properties

This section provides some statistical properties of the OBP-G family of distributions, such as moments, the moment generating function, and entropy.

3.1. Moments

Suppose that X follows the OBP-G family, the r^{th} moments of X is obtained as

$$E(X^r) = \int_{-\infty}^{\infty} x^r f(x) dx, \tag{9}$$

where $f(x)$ is defined in equation (8).

Substituting (8) into (9) gives the moments of the OBP-G as

$$E(X^r) = \frac{1}{B(c, d)} \sum_{i,j=0}^{\infty} \Psi_{i,j} \int_{-\infty}^{\infty} x^r g(x; \varepsilon) (G(x; \varepsilon))^{c+i+j-1} dx.$$

3.2. Moment Generating Function

Assume that a random variable X follows the OBP-G family, the moment generating function of X is given as

$$M_X(t) = E(e^{tx}) = \int_{-\infty}^{\infty} e^{tx} f(x) dx, \tag{10}$$

Inserting (8) in (10), we have

$$M_X(t) = \frac{1}{B(c, d)} \sum_{i,j=0}^{\infty} \Psi_{i,j} \int_{-\infty}^{\infty} e^{tx} g(x; \varepsilon) (G(x; \varepsilon))^{c+i+j-1} dx. \tag{11}$$

Which can be expressed as

$$M_X(t) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} E(X^k). \tag{12}$$

3.3. Entropy

Assume that X is a random variable that follows the OBP-G family, the Rényi entropy [9] of X is expressed as

$$R_{\alpha}(x) = \frac{1}{1-\alpha} \log \left[\int_{-\infty}^{\infty} f^{\alpha}(x) dx \right]; \quad \alpha > 0, \alpha \neq 1, x \in \mathfrak{R}. \tag{13}$$

The integrand $f^{\alpha}(x)$ can be obtained as

$$f^\alpha(x) = [f(x)]^\alpha = \left\{ \frac{g(x; \varepsilon)}{B(c, d)[1-G(x; \varepsilon)]^2} \frac{\left(\frac{G(x; \varepsilon)}{1-G(x; \varepsilon)}\right)^{c-1}}{\left[1 + \left(\frac{G(x; \varepsilon)}{1-G(x; \varepsilon)}\right)\right]^{c+d}} \right\}^\alpha. \quad (14)$$

Therefore, (14) can be rewritten as

$$f^\alpha(x) = \frac{g^\alpha(x; \varepsilon)}{B^\alpha(c, d)[1-G(x; \varepsilon)]^{2\alpha}} \frac{\left(\frac{G(x; \varepsilon)}{1-G(x; \varepsilon)}\right)^{\alpha(c-1)}}{\left[1 + \left(\frac{G(x; \varepsilon)}{1-G(x; \varepsilon)}\right)\right]^{\alpha(c+d)}}. \quad (15)$$

Substituting (5) into (15), we have

$$f^\alpha(x) = \frac{g^\alpha(x; \varepsilon)}{B^\alpha(c, d)[1-G(x; \varepsilon)]^{2\alpha}} \left(\frac{G(x; \varepsilon)}{1-G(x; \varepsilon)}\right)^{\alpha(c-1)} \sum_{l=0}^{\infty} (-1)^l \binom{\alpha(c+d)+l-1}{l} \left(\frac{G(x; \varepsilon)}{1-G(x; \varepsilon)}\right)^l. \quad (16)$$

4. The Odd Beta Prime-Exponential Distribution

This section develops a new probability distribution referred to as the odd beta prime-exponential (OBPE) distribution as a sub-model of the proposed family.

Let X be a random variable with exponential distribution, the cdf and pdf are, respectively, as follows:

$$G(x) = 1 - e^{-bx}, \quad x \geq 0, \quad (17)$$

$$g(x) = be^{-bx}, \quad x \geq 0, \quad (18)$$

where $b > 0$ is the rate parameter.

Hence, the cdf and pdf of the OBPE distribution can be obtained by inserting (17) and (18) in (3) and (4), respectively, as follows:

$$F(x) = \frac{B_{\frac{1-e^{-bx}}{e^{-bx}}}(c, d)}{B(c, d)}, \quad x \geq 0. \quad (19)$$

$$f(x) = \frac{b(1-e^{-bx})^{c-1}}{B(c, d)e^{-bcx} \left[1 + \left(\frac{1-e^{-bx}}{e^{-bx}}\right)\right]^{c+d}}, \quad x \geq 0. \quad (20)$$

5. Applications

In this section, we analyzed two real data sets involving engineering and environment to evaluate the applicability of the OBPE distribution.

4.1. The Airborne Communications Transceiver Data

This engineering data was discussed in [10], and it represents the repair times of 46 failures in (hours) of an airborne communications transceiver.

Here, we will compare the fits of the OBPE with the gamma-exponentiated exponential (GEE) in [11] and the beta-exponential (BE) in [12].

We considered the following criteria to compare these distributions: the values of the negative log-likelihood $(-\hat{\ell})$, Akaike information criteria (AIC), Bayesian information criteria (BIC), Cramer-von Mises (CM), and Anderson-Darling (AD). The smaller the values of these statistics, the better the fit to the data.

The maximum likelihood estimates (MLEs), standard errors (SEs), $-\hat{\ell}$, AIC, BIC, CM, and AD statistics for the OBPE, GEE, and BE are presented in Table 1. From the results in Table 1, it is obvious that the OBPE distribution provides better fits for the data, having smallest values of $-\hat{\ell}$, AIC, BIC, CM, and AD and could be selected as a more appropriate model than other models. Figure 1 depicts the estimated pdfs and cdfs of the fitted distributions. It is obvious from these plots that the OBPE better describes the data than other competing models.

Table 1. MLEs with corresponding SEs (in parentheses), and some statistical measures of competing models for the airborne communications transceiver data.

Distribution	MLE and SE in ()			$-\hat{\ell}$	AIC	BIC	CM	AD
OBPE	$c = 0.6583$ (0.1625)	$d = 1.1018$ (0.1149)	$b = 1.4863$ (4.5831)	100.0163	204.032	207.689	0.054	0.337
GEE	$\alpha = 0.9323$ (0.1701)	$\theta = 0.2585$ (0.0615)	$\lambda = 0.3685$ (0.7650)	104.9309	213.861	217.519	0.175	1.103
BE	$a = 2.6732$ (0.4920)	$b = 2.0190$ (0.2613)	$\lambda = 1.1455$ (0.4016)	128.48	260.960	264.617	0.409	3.007




Figure 1. The estimated pdfs of the OBPE and other competing models for the airborne communications transceiver data.

4.2. Exceedances of Wheaton River Flood Data

This environmental data was analyzed by [13], and it represents the exceedances of flood peaks (in m3/s) of the Wheaton River near Carcross in Yukon Territory, Canada. The data consist of 72 exceedances for the years 1958–1984, rounded to one decimal place.

Also, from the results in Table 2 and the illustrations in Figure 2, it is obvious from these plots that the OBPE provides a better fit to this data than other competing fitted models.

Table 2. MLEs with corresponding SEs (in parentheses), and some statistical measures of competing models for the exceedances of Wheaton River flood data.

Distribution	MLE and SE in ()			$-\hat{\ell}$	AIC	BIC	CM	AD
OBPE	$c = 1.2126$ (0.1166)	$d = 6.7584$ (1.1659)	$b = 1.6077$ (0.3743)	257.839	519.6782	524.231	0.232	1.472
GEE	$\alpha = 10.6378$ (1.3609)	$\theta = 6.6058$ (0.6475)	$\lambda = 2.9650$ (0.4475)	279.958	563.9169	568.470	0.330	2.478
BE	$a = 12.2041$ (1.4391)	$b = 12.2115$ (1.0176)	$\lambda = 5.1753$ (0.2352)	282.334	568.669	573.223	0.469	2.891




Figure 2. The estimated pdfs of the OBPE and other competing models for the exceedances of Wheaton River flood data.

6. Conclusion

A new family of life distributions, called the odd beta prime-G family, is introduced. Some statistical properties of the new family, including moments, the moment generating function, and entropy, are derived. A special sub-model of the new proposed family, called odd beta prime-exponential distribution, is developed, and two real applications are analyzed to demonstrate the flexibility of the new distribution. Empirically, it is proved that the proposed model can give better fits to modeling data than the other competing life distributions.

Author Contributions: Conceptualization, A.A.S., M.O. and H.D.; Methodology, A.A.S., M.O., A.I.I. and M.L.A.; Software, A.A.S. and A.I.I.; Validation, A.A.S., M.O., R.I., H.D., R.I. and A.H.; Supervision, M.O. and H.D.; Formal analysis, A.A.S., M.O., A.I.I., and M.L.A. All authors have read and agreed to the published of the manuscript.

Acknowledgments: The authors would like to thank the Universiti Teknologi PETRONAS for providing support to this project.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- [1] A. Alzaatreh, C. Lee, and F. Famoye, "A new method for generating families of continuous distributions," *Metron*, vol. 71, no. 1, pp. 63-79, 2013.
- [2] M. A. Aljarrah, C. Lee, and F. Famoye, "On generating TX family of distributions using quantile functions," *Journal of Statistical Distributions and Applications*, vol. 1, no. 1, pp. 1-17, 2014.
- [3] N. Eugene, C. Lee, and F. Famoye, "Beta-normal distribution and its applications," *Communications in Statistics-Theory and methods*, vol. 31, no. 4, pp. 497-512, 2002.
- [4] G. M. Cordeiro and M. de Castro, "A new family of generalized distributions," *Journal of statistical computation and simulation*, vol. 81, no. 7, pp. 883-898, 2011.
- [5] H. Torabi and N. H. Montazeri, "The logistic-uniform distribution and its applications," *Communications in Statistics-Simulation and Computation*, vol. 43, no. 10, pp. 2551-2569, 2014.
- [6] M. Bourguignon, R. B. Silva, and G. M. Cordeiro, "The Weibull-G family of probability distributions," *Journal of data science*, vol. 12, no. 1, pp. 53-68, 2014.
- [7] A. Z. Afify, G. M. Cordeiro, M. E. Maed, M. Alizadeh, H. Al-Mofleh, and Z. M. Nofal, "The generalized odd Lindley-G family: Properties and applications," *Anais da Academia Brasileira de Ciências*, vol. 91, 2019.
- [8] A. I. Ishaq and A. A. Abiodun, "The Maxwell-Weibull distribution in modeling lifetime datasets," *Annals of Data Science*, vol. 7, no. 4, pp. 639-662, 2020.
- [9] A. Rényi, "On measures of entropy and information," in *Proceedings of the fourth Berkeley symposium on mathematical statistics and probability*, 1961, vol. 1, no. 547-561: Berkeley, California, USA.

-
- [10] M. K. Refaie, "Burr X exponentiated exponential distribution," *Journal of Statistics and Applications*, vol. 1, no. 2, pp. 71-88, 2018.
- [11] M. M. Ristić and N. Balakrishnan, "The gamma-exponentiated exponential distribution," *Journal of statistical computation and simulation*, vol. 82, no. 8, pp. 1191-1206, 2012.
- [12] S. Nadarajah and S. Kotz, "The beta exponential distribution," *Reliability engineering & system safety*, vol. 91, no. 6, pp. 689-697, 2006.
- [13] V. Choulakian and M. A. Stephens, "Goodness-of-fit tests for the generalized Pareto distribution," *Technometrics*, vol. 43, no. 4, pp. 478-484, 2001.

The Odd Beta Prime-G Family of Probability Distributions.pdf

ORIGINALITY REPORT

24%

SIMILARITY INDEX

17%

INTERNET SOURCES

18%

PUBLICATIONS

13%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

3%

★ Nurul Nnadiyah Zakaria, Mahmud Othman, Rajalingam Sokkalingam, Hanita Daud, Lazim Abdullah, Evizal Abdul Kadir. "Markov Chain Model Development for Forecasting Air Pollution Index of Miri, Sarawak", Sustainability, 2019

Publication

Exclude quotes Off

Exclude matches < 1%

Exclude bibliography On